

Spectral asymptotics of Robin Laplacians on polygonal domains

Magda Khalile

Laboratoire de Mathématiques d'Orsay, Univ. Paris-Sud,
91405 Orsay, France

Abstract

Let $\Omega \subset \mathbb{R}^2$ be a curvilinear polygon and Q_Ω^γ be the Laplacian in $L^2(\Omega)$, $Q_\Omega^\gamma \psi = -\Delta \psi$, with the Robin boundary condition $\partial_\nu \psi = \gamma \psi$, where ∂_ν is the outer normal derivative and $\gamma > 0$. We are interested in the behavior of the eigenvalues of Q_Ω^γ as γ becomes large. We prove that there exists $N_\Omega \in \mathbb{N}$ such that the asymptotics of the N_Ω first eigenvalues of Q_Ω^γ is determined at the leading order by those of model operators associated with the vertices: the Robin Laplacians acting on the tangent sectors associated with $\partial\Omega$. In the particular case of a polygon with straight edges the N_Ω first eigenpairs are exponentially close to those of the model operators. Moreover, if the polygon admits only *non-resonant* or concave corners, we prove that, for any fixed $j \in \mathbb{N}$, the $N_\Omega + j$ eigenvalue $E_{N_\Omega+j}(Q_\Omega^\gamma)$ behaves as

$$E_{N_\Omega+j}(Q_\Omega^\gamma) = -\gamma^2 + \mu_j^D + o(1), \text{ as } \gamma \rightarrow +\infty,$$

where μ_j^D stands for the j th eigenvalue of the operator $D_1 \oplus \dots \oplus D_M$ and D_n denotes the one-dimensional Laplacian $f \mapsto -f''$ on $(0, l_n)$, where l_n is the length of the n th side of Ω , with the Dirichlet boundary condition.